


**Kentucky Annual Tobacco Control Plan, 2002-2003  
Smoking During Pregnancy 1996-2000  
Cumberland Valley District Health Departments**

County	1996		1997		1998		1999		2000	
	N	%	N	%	N	%	N	%	N	%
Bell	435	33.3	384	29.7	403	28.5	357	27.7	371	33.2
Clay	334	41.3	297	41.1	299	30.4	299	37.5	284	42.3
Harlan	439	34.4	444	30.4	421	31.1	438	33.3	408	33.8
Jackson	152	35.5	159	37.1	174	36.8	182	35.2	191	33
Rockcastle	167	32.9	162	35.8	202	26.2	190	28.4	212	26.9
<b>Cumberland Valley</b>	<b>1527</b>	<b>35.6</b>	<b>1446</b>	<b>33.7</b>	<b>1499</b>	<b>30.3</b>	<b>1466</b>	<b>32.4</b>	<b>1466</b>	<b>34.2</b>
<b>Kentucky</b>	<b>50264</b>	<b>24.5</b>	<b>50338</b>	<b>24.7</b>	<b>54125</b>	<b>23.7</b>	<b>53216</b>	<b>23.7</b>	<b>55978</b>	<b>23.4</b>



Healthy Kentuckians 2010: Prevention Initiative

Objective 3.4. Reduce cigarette smoking among pregnant women to a prevalence of no more than 17 percent. (1997 Baseline: 25 percent).

Objective 3.5. Increase to at least 50 percent the proportion of pregnant women who abstain from tobacco use beginning early in pregnancy and maintain abstinence for the remainder of their pregnancy, following delivery, and through 6 weeks postpartum.

Sources. Healthy Kentuckians 2010 Spring 2000  
Kentucky Vital Statistics Birth File 1996-00