


**Kentucky Annual Tobacco Control Plan, 2002-2003  
Smoking During Pregnancy 1996-2000  
Green River Health Department**

County	1996		1997		1998		1999		2000	
	N	%	N	%	N	%	N	%	N	%
Daviess	1195	24.4	1242	28.0	1339	26.1	1282	23.4	1289	22.4
Hancock	114	21.1	122	24.6	110	23.6	101	26.7	153	23.5
Henderson	491	15.9	466	20.8	562	20.5	509	18.5	614	16.8
McLean	110	34.5	110	24.5	130	23.1	130	27.7	155	27.1
Ohio	252	34.5	275	30.5	286	35.0	294	25.9	314	22.3
Union	189	9.5	180	17.2	179	17.3	165	13.3	200	15.5
Webster	168	14.3	164	20.1	181	23.8	164	23.8	189	21.2
<b>Green River</b>	<b>2519</b>	<b>22.2</b>	<b>2559</b>	<b>25.4</b>	<b>2787</b>	<b>24.9</b>	<b>2645</b>	<b>22.5</b>	<b>2914</b>	<b>21</b>
<b>Kentucky</b>	<b>50264</b>	<b>24.5</b>	<b>50338</b>	<b>24.7</b>	<b>54125</b>	<b>23.7</b>	<b>53216</b>	<b>23.7</b>	<b>55978</b>	<b>23.4</b>



Healthy Kentuckians 2010: Prevention Initiative

Objective 3.4. Reduce cigarette smoking among pregnant women to a prevalence of no more than 17 percent. (1997 Baseline: 25 percent).

Objective 3.5. Increase to at least 50 percent the proportion of pregnant women who abstain from tobacco use beginning early in pregnancy and maintain abstinence for the remainder of their pregnancy, following delivery, and through 6 weeks postpartum.

Sources. Healthy Kentuckians 2010 Spring 2000  
Kentucky Vital Statistics Birth File 1996-00