# KENTUCKY LOCAL HEALTH DEPARTMENT TOBACCO CESSATION SURVEY CHRISTIAN COUNTY HEALTH DEPARTMENT JUNE-AUGUST 2000

#### Do you or other health department staff provide smoking cessation programs?

| Christian | Kentucky ( <u>n</u> =54) |
|-----------|--------------------------|
| No        | 29(53.7%)                |

### If "yes," what programs do your health department provide?

|                              | Christian | Kentucky ( <u>n</u> =29) |
|------------------------------|-----------|--------------------------|
| Fresh Start                  |           | 15(51.7%)                |
| Freedom from Smoking         |           | 1(3.4%)                  |
| Cooper-Clayton Method        |           | 24(82.8%)                |
| Nicotine Replacement Therapy |           | 7(24.1%)                 |

#### If "yes," how many smoking cessation programs were offered within the last year?

| Christian | Kentucky ( <u>n</u> =26) |
|-----------|--------------------------|
|           | 3.73±2.65, range=0-12    |

### If "yes," how many clients/patients participated in smoking cessation programs within the last year?

| Christian | Kentucky ( <u>n</u> =26) |
|-----------|--------------------------|
|           | 33.27±47.23, range=0-227 |

#### If "yes," how many clients/patients completed smoking cessation programs within the last year?

| Christian | Kentucky ( <u>n</u> =25) |
|-----------|--------------------------|
|           | 18.08±23.93, range=0-105 |

#### If "yes," what is the job title of the health department staff providing smoking cessation?

|                 | Christian | Kentucky ( <u>n</u> =29) |
|-----------------|-----------|--------------------------|
| Health Educator |           | 26(89.7%)                |
| Nurse           |           | 11(37.9%)                |
| Other           |           | 6(20.7%)                 |

On average, how many requests for smoking cessation does your health department receive per month?

| Christian | Kentucky ( <u>n</u> =54) |
|-----------|--------------------------|
| 0         | 4.63±8.51, range=0-50    |

Within the past year, have any 'Cooper-Clayton Method to Stop Smoking' programs been provided in your county?

| Christian | Kentucky ( <u>n</u> =54) |
|-----------|--------------------------|
| No        | 23(42.6%)                |

If "yes," how many people are trained to provide Cooper-Clayton programs?

| Christian | Kentucky ( <u>n</u> =21) |
|-----------|--------------------------|
|           | 3.10±3.35, range=0-13    |

Within the past year, has your health department provided educational programs on smoking cessation for health care providers?

| Christian | Kentucky ( <u>n</u> =53) |
|-----------|--------------------------|
| No        | 8(15.1%)                 |

If "yes," about how many programs did you provide?

| Christian | Kentucky ( <u>n</u> =6) |
|-----------|-------------------------|
|           | 2.33±1.63, range=0-4    |

Does your health department use a tobacco-user identification system for every patient at every clinic visit?

| Christian | Kentucky ( <u>n</u> =53) |
|-----------|--------------------------|
| Yes       | 44(83.0%)                |

#### What kinds of educational methods are used for tobacco use cessation in your health department?

|                       | Christian | Kentucky ( <u>n</u> =54) |
|-----------------------|-----------|--------------------------|
| Formal group classes  |           | 19(35.2%)                |
| Individual counseling | ✓         | 50(92.6%)                |
| Audiovisual materials | ✓         | 25(46.3%)                |
| Printed materials     | ✓         | 50(92.6%)                |
| Posters               | ✓         | 45(83.3%)                |
| Other                 |           |                          |

#### Does your clinic have self-help materials for tobacco use cessation available in every exam room?

| Christian | Kentucky ( <u>n</u> =54) |
|-----------|--------------------------|
| No        | 20(37.0%)                |

#### Are materials on secondhand smoke available in every exam room?

| Christian | Kentucky ( <u>n</u> =54) |
|-----------|--------------------------|
| No        | 21(39.0%)                |

#### Do your health care providers routinely:

|  | Christian | Kentucky     |
|--|-----------|--------------|
| Ask patients about secondhand smoke exposure in their homes? | Yes       | 43/53(81.1%) |
| Advise patients to ban smoking in their homes?               | Yes       | 45/53(84.9%) |
| Advise patients to ban smoking in their cars?                | Yes       | 35/50(70.0%) |

### Do your health care providers routinely refer tobacco users who want to quit to a counselor or to a program available in your community for tobacco use cessation treatment?

| Christian | Kentucky (n=54) |
|-----------|-----------------|
| Yes       | 41(76.0%)       |

#### Does your employee health plan reimburse for tobacco use cessation treatment?

| Christian | Kentucky (n=43) |
|-----------|-----------------|
| No        | 3(7.0%)         |

## How frequently do your health care providers prescribe or recommend the following medications for tobacco use cessation? (those that prescribe often or very often)

|                      | Christian | Kentucky    |
|----------------------|-----------|-------------|
| Nicotine Gum         |           | 6/50(12%)   |
| Nicotine Patch       |           | 11/50(22%)  |
| Nicotine Nasal Spray |           | 0/51(0%)    |
| Nicotine Inhaler     |           | 0/51(0%)    |
| Bupropion            |           | 8/51(15.7%) |
| Clonidine            |           | 0/51(0%)    |
| Nortriptyline        |           | 1/51(2%)    |

#### What are the obstacles to providing smoking cessation services? (those that agree or strongly agree)

|                   | Christian | Kentucky     |
|-------------------|-----------|--------------|
| Not enough staff  | ✓         | 25/54(46.3%) |
| Staff not trained | ✓         | 28/54(51.9%) |
| Too busy          | ✓         | 12/54(22.2%) |
| Few requests      | ✓         | 34/54(63.0%) |
| Too expensive     | ✓         | 24/51(47.1%) |
| Ineffective       |           | 16/53(30.2%) |

Data compiled by Dr. Ellen J. Hahn and Dr. Mary Kay Rayens, University of Kentucky College of Nursing, Lexington, KY 40536-0232, (859) 257-2358, ejhahn00@pop.uky.edu