KENTUCKY LOCAL HEALTH DEPARTMENT TOBACCO CESSATION SURVEY PENNYRILE DISTRICT HEALTH DEPARTMENT JUNE-AUGUST 2000

Do you or other health department staff provide smoking cessation programs?

| Pennyrile | Kentucky (<u>n</u> =54) |
|-----------|--------------------------|
| No | 29(53.7%) |

If "yes," what programs do your health department provide?

| | Pennyrile | Kentucky (<u>n</u> =29) |
|------------------------------|-----------|--------------------------|
| Fresh Start | | 15(51.7%) |
| Freedom from Smoking | | 1(3.4%) |
| Cooper-Clayton Method | | 24(82.8%) |
| Nicotine Replacement Therapy | | 7(24.1%) |

If "yes," how many smoking cessation programs were offered within the last year?

| Pennyrile | Kentucky (<u>n</u> =26) |
|-----------|--------------------------|
| | 3.73±2.65, range=0-12 |

If "yes," how many clients/patients participated in smoking cessation programs within the last year?

| Pennyrile | Kentucky (<u>n</u> =26) |
|-----------|--------------------------|
| | 33.27±47.23, range=0-227 |

If "yes," how many clients/patients completed smoking cessation programs within the last year?

| Pennyrile | Kentucky (<u>n</u> =25) |
|-----------|--------------------------|
| | 18.08±23.93, range=0-105 |

If "yes," what is the job title of the health department staff providing smoking cessation?

| | Pennyrile | Kentucky (<u>n</u> =29) |
|-----------------|-----------|--------------------------|
| Health Educator | | 26(89.7%) |
| Nurse | | 11(37.9%) |
| Other | | 6(20.7%) |

On average, how many requests for smoking cessation does your health department receive per month?

| Pennyrile | Kentucky (<u>n</u> =54) |
|-----------|--------------------------|
| 0 | 4.63±8.51, range=0-50 |

Within the past year, have any 'Cooper-Clayton Method to Stop Smoking' programs been provided in your county?

| Pennyrile | Kentucky (<u>n</u> =54) |
|-----------|--------------------------|
| No | 23(42.6%) |

If "yes," how many people are trained to provide Cooper-Clayton programs?

| Pennyrile | Kentucky (<u>n</u> =21) |
|-----------|--------------------------|
| | 3.10±3.35, range=0-13 |

Within the past year, has your health department provided educational programs on smoking cessation for health care providers?

| Pennyrile | Kentucky (<u>n</u> =53) |
|-----------|--------------------------|
| No | 8(15.1%) |

If "yes," about how many programs did you provide?

| Pennyrile | Kentucky (<u>n</u> =6) |
|-----------|-------------------------|
| | 2.33±1.63, range=0-4 |

Does your health department use a tobacco-user identification system for every patient at every clinic visit?

| Pennyrile | Kentucky (<u>n</u> =53) |
|-----------|--------------------------|
| Yes | 44(83.0%) |

What kinds of educational methods are used for tobacco use cessation in your health department?

| | Pennyrile | Kentucky (<u>n</u> =54) |
|-----------------------|-----------|--------------------------|
| Formal group classes | ✓ | 19(35.2%) |
| Individual counseling | ✓ | 50(92.6%) |
| Audiovisual materials | ✓ | 25(46.3%) |
| Printed materials | ✓ | 50(92.6%) |
| Posters | ✓ | 45(83.3%) |
| Other | | |

Does your clinic have self-help materials for tobacco use cessation available in every exam room?

| Pennyrile | Kentucky (<u>n</u> =54) |
|-----------|--------------------------|
| No | 20(37.0%) |

Are materials on secondhand smoke available in every exam room?

| Pennyrile | Kentucky (<u>n</u> =54) |
|-----------|--------------------------|
| No | 21(39.0%) |

Do your health care providers routinely:

| | Pennyrile | Kentucky |
|--|-----------|--------------|
| Ask patients about secondhand smoke exposure in their homes? | Yes | 43/53(81.1%) |
| Advise patients to ban smoking in their homes? | Yes | 45/53(84.9%) |
| Advise patients to ban smoking in their cars? | No | 35/50(70.0%) |

Do your health care providers routinely refer tobacco users who want to quit to a counselor or to a program available in your community for tobacco use cessation treatment?

| Pennyrile | Kentucky (n=54) |
|-----------|-----------------|
| No | 41(76.0%) |

Does your employee health plan reimburse for tobacco use cessation treatment?

| Pennyrile | Kentucky (n=43) |
|-----------|-----------------|
| No | 3(7.0%) |

How frequently do your health care providers prescribe or recommend the following medications for tobacco use cessation? (those that prescribe often or very often)

| | Pennyrile | Kentucky |
|----------------------|-----------|-------------|
| Nicotine Gum | | 6/50(12%) |
| Nicotine Patch | | 11/50(22%) |
| Nicotine Nasal Spray | | 0/51(0%) |
| Nicotine Inhaler | | 0/51(0%) |
| Bupropion | | 8/51(15.7%) |
| Clonidine | | 0/51(0%) |
| Nortriptyline | | 1/51(2%) |

What are the obstacles to providing smoking cessation services? (those that agree or strongly agree)

| | Pennyrile | Kentucky |
|-------------------|-----------|--------------|
| Not enough staff | ✓ | 25/54(46.3%) |
| Staff not trained | ✓ | 28/54(51.9%) |
| Too busy | | 12/54(22.2%) |
| Few requests | ✓ | 34/54(63.0%) |
| Too expensive | | 24/51(47.1%) |
| Ineffective | ✓ | 16/53(30.2%) |

Data compiled by Dr. Ellen J. Hahn and Dr. Mary Kay Rayens, University of Kentucky College of Nursing, Lexington, KY 40536-0232, (859) 257-2358, ejhahn00@pop.uky.edu