North Central District Health Department Tobacco Cessation Survey, 2003

		North Central	Kentucky (N=55)
1.	Are you or other health department staff involved in providing tobacco cessation programs?	Yes	55 (100.0%) Yes
2.	If "Yes," what programs do you provide?		
	a. Fresh Start (American Cancer Society)	No	6 (10.9%) Yes
	b. Freedom from Smoking (American Lung Association)	No	0 (0%) Yes
	c. Cooper-Clayton Method to Stop Smoking	Yes	51 (92.7%) Yes
	d. Make Yours a Fresh Start Family (American Cancer Society)	Yes	51 (92.7%) Yes
	e. Quit and Win	No	25 (45.5%) Yes
	f. TEG/TAP	Yes	42 (76.4%) Yes
	g. NOT	No	8 (14.5%) Yes
3.	If "Yes," how many tobacco cessation programs were offered within the last year?	6	Total Programs = 504 Mean = 9.5 ($\underline{SD} = 12.8$)
4.	If "Yes," how many clients/patients participated in tobacco cessation programs within the last year?	76 (81 ^a)	Total Clients/Patients = 4439 (5277 ^a) Mean = 83.8 (\underline{SD} = 87.0)
5.	If "Yes," how many clients/patients completed tobacco cessation programs within the last year?	74	Total Clients/Patients = 2580 Mean = 49.6 (SD = 50.4) (58.1% Completion Rate)
6.	If "Yes," what is the job title of health department staff providing tobacco cessation?		
	a. Health Educator	Yes	37 (67.3%) Yes
	b. Nurse	Yes	43 (78.2%) Yes
	c. Social Worker	No	7 (12.7%) Yes
	d. Physician	No	0 (0%) Yes

^aNumber of participants in LHD cessation programs + number of Quit and Win registrants - number enrolled in both (see Question 8).

The Kentucky Department for Public Health

University of Kentucky College of Nursing North Central District Health Department Tobacco Cessation Survey, 2003

		North Central	Kentucky (N=55)
7.	On average, how many requests for tobacco cessation does your health department receive per month?	2	Total Requests = 455 Mean = $9.10 (\underline{SD} = 16.5)$
8.	According to our records, 5 person in your area participated in the Quit & Win Tobacco-Free Contest. To the best of your knowledge, about how many people involved in your cessation programs also participated in Quit and Win 2003?	0	Total People = 137 Mean = 5.07 (\underline{SD} = 10.9)
9.	Within the past year, have any Cooper-Clayton Method to Stop Smoking programs been provided in your service area?	Yes	47 (85.5%) Yes
10.	If "Yes," how many people in your service area are trained to provide Cooper-Clayton programs?	2	Total People = 290 Mean = 6.4 (\underline{SD} = 7.7)
11.	How are the tobacco cessation programs in your area promoted?		
	a. Word of mouth	Yes	55 (100%) Yes
	b. Flyers/Brochures	Yes	53 (96.4%) Yes
	c. Newspapers	Yes	52 (94.5%) Yes
	d. Radio	No	35 (63.6%) Yes
	e. Television	No	18 (32.7%) Yes
	f. Church or Community bulletins	No	30 (54.5%) Yes
	g. No promotion	No	0 (0%) Yes
12.	Approximately how much money have you spent in the past year promoting your cessation programs through newspaper, radio or television?	\$200.00	Total Amount =\$163,854.00 Mean =\$3,486.30 (<u>SD</u> = \$9061.36)
13.	Within the past year, has your health department provided education programs on tobacco cessation for health care providers?	Yes	19 (35.2%) Yes
14.	If "Yes," about how many programs did you provide?	1	Total Programs = 39 Mean = $2.05 (\underline{SD} = 1.7)$

North Central District Health Department Tobacco Cessation Survey, 2003

		North Central	Kentucky (N=55)
15.	Does your health department have a tobacco-user identification system (e.g. tobacco use as a vital sign, flagging the charts)?	Yes	48 (87.3%) Yes
16.	If "Yes," is the tobacco-user identification system used for every patient at every clinic visit?	Yes	44 (91.7%) Yes
17.	What kinds of educational methods are used for tobacco use cessation in your health department?		
	a. Formal group classes	Yes	42 (76.4%) Yes
	b. Individual counseling	Yes	54 (98.2%) Yes
	c. Audio-visual materials	No	45 (81.8%) Yes
	d. Printed materials	Yes	55 (100.0%) Yes
	e. Posters	Yes	53 (96.4%) Yes
18.	Does your clinic have self-help materials for tobacco use cessation available in every exam room?	Yes	43 (78.2%) Yes
19.	Are materials on secondhand smoke available in every exam room?	Yes	37 (67.3%) Yes
20.	Do your health care providers routinely ask patients about secondhand smoke exposure in the home?	Yes	50 (90.9%) Yes
21.	Do your health care providers routinely advise patients to ban smoking in the home?	Yes	51 (94.4%) Yes
22.	Do you health care providers routinely advise patients to ban smoking in their cars?	Yes	51 (94.4%) Yes
23.	Do your health care providers routinely provide brief cessation interventions using the 5 A's (Ask, Advise, Assess, Assist, Arrange)?	Yes	35 (64.8%) Yes
24.	Do your providers routinely refer tobacco users who want to quit to a counselor or to a program available in your community for tobacco cessation treatment?	Yes	53 (98.1%) Yes

^{*=} missing data

The Kentucky Department for Public Health University of Kentucky College of Nursing North Central District Health Department Tobacco Cessation Survey, 2003

		North Central	Kentucky (N=55)	
25.	Does your employee health plan(s) at your health department reimburse for tobacco use cessation treatment? (including nicotine gum/patch and/or counseling)	No	24 (47.1%) Yes	
26.	How frequently do your health care providers prescribe or recommend the following medications for tobacco use cessation?			
	a. Nicotine Gum	2=	1= Never 2= 3= 4= 5= Very Often	40.0% 16.4% 14.5% 9.1% 20.0%
	b. Nicotine Patch	2=	1= Never 2= 3= 4= 5= Very Often	32.7% 5.5% 9.1% 20.0% 32.7%
	c. Nicotine Nasal Spray	1= Never	1= Never 2= 3= 4= 5= Very Often	64.8% 22.2% 11.1% 0.0% 1.9%
	d. Nicotine Inhaler	1= Never	1= Never 2= 3= 4= 5= Very Often	68.5% 20.4% 5.6% 1.9% 3.7%
	e. Nicotine Lozenges	1= Never	1= Never 2= 3= 4= 5= Very Often	70.4% 11.1% 9.3% 3.7% 5.6%

North Central District Health Department Tobacco Cessation Survey, 2003

	·	North Central	Kentucky (N=55)	
	f. Buproprion (Wellbutrin, Zyban)	1= Never	1= Never	49.1%
			2=	12.7%
			3=	12.7%
			4=	16.4%
			5= Very Often	9.1%
	g. Clonidine (suppresses opoid withdrawal symptoms;	1= Never	1= Never	78.8%
	typically used to treat high blood pressure)		2=	9.6%
			3=	11.5%
			4=	0.0%
			5= Very Often	0.0%
	h. Nortriptyline (antidepressant)	1= Never	1= Never	80.8%
			2=	9.6%
			3=	9.6%
			4=	0.0%
			5= Very Often	0.0%
27.	We have enough staff to provide tobacco cessation programs.	1= Strongly Disagree	1= Strongly Disagree	16.4%
			2= Disagree	21.8%
			3= Agree	40.0%
			4= Strongly Agree	21.8%
28.	Our staff are not trained to provide tobacco cessation.	3= Agree	1= Strongly Disagree	36.4%
			2= Disagree	43.6%
			3= Agree	14.5%
			4= Strongly Agree	5.5%
29.	We are too busy to provide tobacco cessation.	3= Agree	1= Strongly Disagree	30.9%
			2= Disagree	52.7%
			3= Agree	16.4%
			4= Strongly Agree	0.0%
30.	We have few requests for tobacco cessation programs.	3= Agree	1= Strongly Disagree	24.1%
			2= Disagree	31.5%
			3= Agree	29.6%
			4= Strongly Agree	14.8%

North Central District Health Department Tobacco Cessation Survey, 2003

		North Central	Kentucky (N=55)	
31.	Tobacco cessation programs are too expensive.	2= Disagree	1= Strongly Disagree 2= Disagree 3= Agree 4= Strongly Agree	38.2% 41.8% 14.5% 5.5%
32.	Tobacco cessation programs are effective.	3= Agree	1= Strongly Disagree 2= Disagree 3= Agree 4= Strongly Agree	0.0% 1.9% 64.2% 34.0%

This report was prepared through a contract between the Kentucky Cabinet for Health Services and the University of Kentucky College of Nursing

For more information, contact:
Dr. Ellen Hahn, DNS, RN
Associate Professor
University of Kentucky
College of Nursing
(859) 257-2358
ejhahn00@pop.uky.edu