


**Kentucky Annual Tobacco Control Plan, 2002-2003
Smoking During Pregnancy 1996-2000
Three Rivers District Health Department**

County	1996		1997		1998		1999		2000	
	N	%	N	%	N	%	N	%	N	%
Carroll	119	34.5	118	36.4	112	36.6	122	32.8	151	37.7
Gallatin	91	41.8	101	31.7	115	31.3	110	39.1	114	35.1
Owen	85	27.1	117	36.8	134	30.6	118	30.5	128	37.5
Pendleton	168	32.7	172	38.4	194	29.4	160	30.6	203	30
Three Rivers	463	33.9	508	36.2	555	31.5	510	32.9	596	34.6
Kentucky	50264	24.5	50338	24.7	54125	23.7	53216	23.7	55978	23.4



Healthy Kentuckians 2010: Prevention Initiative

Objective 3.4. Reduce cigarette smoking among pregnant women to a prevalence of no more than 17 percent. (1997 Baseline: 25 percent).

Objective 3.5. Increase to at least 50 percent the proportion of pregnant women who abstain from tobacco use beginning early in pregnancy and maintain abstinence for the remainder of their pregnancy, following delivery, and through 6 weeks postpartum.

Sources. Healthy Kentuckians 2010 Spring 2000
Kentucky Vital Statistics Birth File 1996-00